

## The Significance of Plasma Sodium

By

**Wandee Varavithya M.D.\***

---

The clinical interpretation of plasma sodium concentration is based on the following premises :

1. Body fluids are constantly in osmotic equilibrium. (1,2) Important exceptions are fluids recently introduced into stomach and fluids in the interstitial spaces of the renal medulla, certain actively secreting glands, and the tubular epithelial cells of various nephron segments.

2. Osmotic equilibrium between the cells and the surrounding fluid is maintained by transfer of water. (3,4). Important exceptions are red blood cells and the cells of central nervous system which may respond to osmotic alterations in the extracellular environment by transfer of both water and solute. (5-7)

3. Change in body water content are reflected by changes in plasma sodium concentration. (8-11) In general, a significant alteration in plasma sodium concentration is an evidence of an abnormality in body water content not in body sodium content.

### Plasma Sodium and Effective Body Fluid Osmolality

The qualitative differences between cellular and extracellular fluids are maintained by the transport and per-

meability characteristics of cell membranes. In general, cell membranes are relatively impermeable to the sodium ion, so sodium and its associated anions are functionally restricted to the extracellular water. Potassium and the anions associated with it are relatively restricted to intracellular water and account for the major fraction of intracellular milliosmols. Intracellular fluid is in osmotic equilibrium with interstitial fluid and plasma, so the osmotic characteristics of plasma may be used to describe the osmolality of most body fluids. Sodium is the most abundant ion in extracellular fluid and accounts for about 90% of the cations in plasma. Most of the anions in extracellular fluid are also univalent, therefore twice the concentration of plasma sodium is a good approximation of 90% of the total ions in plasma. (12) It is fortuitous that the osmotic coefficient of an 0.300 M sodium chloride solution is 0.93, making 90% of the total ion concentration in plasma a good approximation for effective body fluid osmolality. (13)

The major clinical value of plasma sodium is its reliability as a parameter for effective body fluid osmolality. There are two sets of conditions in which concentration of sodium in plasma may

---

\* แผนกกุมารเวชศาสตร์ คณะแพทยศาสตร์ โรงพยาบาลรามาธิบดี.

be lower than normal when body fluid osmolarity is normal or increased. Hyponatremia may occur in patients with hyperproteinemia. (14) The concentration of sodium in water phase of plasma is normal if body fluid osmolarity is normal. Plasma sodium concentration may also be reduced, in the presence of normal or increased body fluid osmolarity, when the extracellular space is expanded by the accumulation of nonionic extracellular solutes such as glucose, or mannitol. (15-18) Excluding these conditions, plasma sodium concentration is a reliable parameter for effective body fluid osmolarity.

### Total Plasma Solute Concentration by Freezing Point Depression

The estimation of total plasma solute concentration by depression of freezing point is now commonly used in both investigative and clinical laboratories. Using freezing point depression, it is necessary to distinguish between total and effective plasma osmolality. The effective plasma osmolality reflects the concentration of plasma solute which influences the distribution of body water. Urea and sugar usually account for 5 to 15 milliosmols of solute per kilogram of plasma water when total plasma solute concentration is estimated by free-

zing point depression. These solute particles generally have little influence on the distribution of body water. Under most circumstances the milliosmols due to urea and glucose should be subtracted from total plasma solute concentration to define the physiologically effective solute.

Observations were made on 150 pediatric patients. These children did not have hyperlipemia, hyperproteinemia, of blood sugar levels above 180 mg%, so plasma was considered a reliable parameter for effective body fluid osmolality. The patients ranged in age from six days to 13 years and had a variety of infectious diseases, metabolic disorders, and congenital anomalies. Dehydration due to gastroenteritis was the most common diagnosis. Plasma sodium concentrations ranged from 133 to 207 mEq/L, with a mean of  $138 \pm 4$  mEq/L. Total plasma osmolarity varied from 235 to 422 mOsm/Kg, with a mean of  $292 \pm 33$ , and effective plasma osmolarity from 215 to 398 mOsm/Kg, with a mean of  $274 \pm 27$  mOsm/Kg. Plasma urea nitrogen concentrations varied from 3 to 268 mg% and plasma sugar from 29 to 176 mg%

Effective plasma osmolarity was derived by subtraction from total plasma solute concentration the milliosmols

contributed by glucose and urea.\*When plots were made between the plasma solute concentration and plasma sodium concentration, and between the latter and effective plasma solute concentration, the correlation between the total plasma osmolarity and plasma sodium concentration is highly significant ( $r=0.89$ ,  $p 0.01$ ). The relation between effective plasma osmolarity and plasma sodium concentration is even closer, ( $r=0.95$ ,  $p 0.01$ ). The difference between the two correlation coefficients is a reflection of the measurement, by freezing point depression, of the physiologically ineffective milliosmols contributed by glucose and urea.

Edelman and his associates showed good correlation between total serum solute concentration and serum sodium in 98 adult patients ( $r = 0.8$ ). (19) Even better correlation was evident in the relationship between effective plasma osmolarity and the concentration of sodium in plasma water. Similar results were reported by Hellerstein, et al in 38 pediatric patients. (7)

Edelman (19)

$$Y = 175x + 10.1, r = 0.97 \text{ (No.=98)}$$

Hellerstein

$$Y = 186x - 2.4, r = 0.96 \text{ (No.=38)}$$

Glucose, Urea, and other nonionic

extracellular solute may affect the distribution of body water. (20) Abrupt increase in the plasma urea concentration causes a transient rise in effective osmolarity. These effects are dissipated in a few hours as the concentration of urea in cell water approaches that in extracellular water. (12) Extracellular glucose undoubtedly exerts effective osmotic activity above certain plasma glucose levels, although the specific threshold is not known. This probably varies between physiological events as well as between individuals. In the total absence of insulin activity, glucose is probably an osmotically effective solute for most cells at rather low plasma levels.

For the estimation of effective osmolarity from total plasma solute concentration, extracellular fluid glucose may be treated as if it does not contribute effective solute at plasma concentration below 180 mg% (10 mOsm/Kg), the approximate renal threshold for glucose.

The normal range for effective plasma osmolarity was  $273 \pm 5.4$  mOsm/Kg plasma water in a group of 77 children ranging in age from one month to fifteen years. The plasma sodium range in this group of children was  $139 \pm 2.9$  mEq/L.

---


$$* [mOsm]_{EFF} = [mOsm]_{FP} - \frac{[Urea \text{ Nitrogen } mg\% \times 10]}{28} + \frac{[Sugar \text{ mg}\% \times 10]}{180}$$

$[mOsm]_{EFF}$  is effective plasma osmolarity and  $[mOsm]_{FP}$  is total plasma solute concentration by freezing point depression. The milliosmols per kilogram of plasma water contributed by sugar and urea assumed to be equal to the sum of the millimolar concentration of these solutes per liter of plasma.

### Summary

In the absence of hyperlipemia, and hyperproteinemia, there is a close correlation between plasma sodium concentration and effective plasma osmolarity. In the clinical interpretation, plasma sodium concentration is a reliable parameter for effective body fluid osmolarity.

### References

1. Gamble, J.L., Ross, G.S. and Tisdall, F.F.; The Metabolism of Fixed Base during, *J. Biol Chem*, 57:633, 1923.
2. Darrow, D.C., and Yannet, H.; The Changes in the Distribution of Body Water Accompany Increase and Decrease in Extracellular Electrolyte, *J. Clin. Invest.* 14: 266, 275, 1935.
3. Wynn, V.: A Metabolic Study of Acute Water Intoxication in Man and Dogs, *Clinical Science* 14: 669, 1955.
4. Leaf, A., Chatillon, J.Y., Wrong, O., and Tuttle, E.P.: The Mechanism of the Osmotic Adjustment of Body Cells as determined in Vivo by the Volume of Distribution of a large Water Load, *J Clin Invest*, 33: 1261, 1954.
5. Yannet, H.: Changes in Brain resulting from Depletion of Extracellular Electrolytes, *Am J Physiol*, 128: 683-689, 1940.
6. Ames, A., Isom, J.B., and Nesbett, F.B.: Effects of Osmotic Changes on Water and Electrolytes in Nervous Tissue, *J Physiol*, 177: 246-262, 1965.
7. Hellerstein, S., Varavitha, W. and Graddy, D.: Plasma and Red Blood Cell and Water Solute, *Am. J Dis Child* 112: 298, 1966
8. Darrow, D.C. and Hellerstein, S.: Interpretation of certain Changes in Body Water and Electrolytes, *Physiol Rev*, 38: 114, 1958.
9. Leaf, A.: The Clinical and Physiological Significant of the Serum Sodium Concentration, *New Eng J Med*, 267: 24-77, 1962.
10. Bartter, F.C.: Regulation of the Volume and Composition of Extracellular and Intracellular fluid, *Ann N.Y. Acad Sci*, 110: 682, 1963.
11. Berliner, R.W.; The Regulation of Water Balance and Plasma Sodium Concentration, *Arch Int Med*, 102: 986, 1958.
12. Pitts, R.F.: Physiology of the Kidney and Body Fluids, Year Book, Chicago, 1963.
13. Wolf, A.V.: Aqueous Solutions and Body Fluids, New York Harper E. Row Publisher Inc, 1966.
14. Albrink, M.J., et al; The Displacement of Serum Water by the Lipids of Hyperlipemic Serum: A new Method for the Rapid Determination of Serum Water, *J Clin Invest*, 34: 1483-1488, 1955.
15. Wright, H.K., and Gann, D.S.: Hyperglycemic Hyponatremia in Non-diabetic Patients, *Arch In Med*, 112: 344-346, 1963.

16. Sunderman, F.W. and Williams, E.S.: Studies in Serum Electrolytes: IX. The Changes in Total Quantity and Osmolal Concentration of Glucose and Chloride in the Serum after the Ingestion of Glucose of Diabetic Patients, *J Clin Invest*, 14: 245-250 1935.

17. Wise, B.L.: Effects of Infusion of Hypertonic Mannitol on Electrolytes and on Osmolarity of Serum and Cerebrospinal Fluid, *J Neuro Surg*, 20: 961, 1963.

18. Edelman, I.S., et al: Interrelations between Serum Osmolarity and total Exchangable Sodium, total Exchangable Potassium and total Body Water, *J Clin Invest*, 37: 1236-1256; 1958.

19. Dodge, P.R., Crawford, L.D., and Brobst, J.H.: Studies in Experimental Water Intoxication, *Arch Neurol*, 5: 513, 1960.

20. Javid, M.: Effect of Urea on Cerebrospinal Fluid Pressure in Human Subjects, *J.A.M.A.* 160: 943-949' 1960.

---

# IMMUNOLOGY

## IN THE FIELD OF OBSTETRICS & GYNECOLOGY

นายแพทย์ ม.ล. ตะวันฉาย ศิริวงศ์\*

ในระหว่าง ๓๐ ปี ที่ล่วงมานี้ นักวิทยาศาสตร์ทางการแพทย์ และวิจัยจำนวนมากมาย ได้พุ่งความสนใจมายัง Immunology ซึ่งเป็นผลทำให้มีการวิจัยทางแขนงวิชาอื่น เพิ่มมากขึ้นทุกที และนักวิจัยกลุ่มหนึ่ง หันความสนใจมาเฉพาะแขนงวิชาอื่นในส่วนที่เกี่ยวข้องกับวิชาสูติศาสตร์-นรีเวชวิทยาโดยเฉพาะ

### การทดสอบการตั้งครรภ์ (Pregnancy test)

เริ่มต้นด้วย Wide และ Gemzell ในปี ค.ศ. ๑๙๖๐ ได้เปิดเผยเทคนิคในการตรวจหา Human Chorionic gonadotropin (HCG)- โดยวิธี Hemagglutination Inhibition ซึ่งปรากฏว่าได้ผลดีเลิศ, รวดเร็ว และราคาถูกลงกว่าการทดสอบการตั้งครรภ์โดยวิธีอื่นๆ

วัสดุ ที่ใช้ในการทดสอบประกอบด้วย Rabbit serum containing antibodies to HCG (ได้จากกระต่ายซึ่งถูกฉีดด้วย Pure

HCG) และ HCG-coated red-cells (ได้จากเม็ดโลหิตแดงของแกะ Stabilized ด้วยฟอร์มาลีน tanned ด้วย tannic acid เพื่อ facilitate การติดเชื่อมต่อ HCG และผสมกับ HCG)

การทดสอบคือ ผสมบัสสาวะของผู้ป่วยด้วยซีรัมของกระต่ายซึ่งมี HCG แอนติบอดีแล้วเติมด้วย HCG coated red cells ถ้าคนไข้ตั้งครรภ์จริง บัสสาวะของคนไข้จะต้องมี HCG- ซึ่งจะทำการปฏิกิริยากับแอนติบอดี ในซีรัมของกระต่าย เพราะฉะนั้นเมื่อใส่ HCG Coated red cells เข้าไปจะไม่มีอาการรวมตัว (agglutinate) และจะตกตะกอน รวมตัวกันอยู่ที่ก้นหลอดทดลอง แต่ถ้าผู้ป่วยไม่ตั้งครรภ์ บัสสาวะของคนไข้รายนั้นจะไม่มี HCG อยู่เลยและ HCG antibodies ในซีรัมของกระต่ายจะคงอยู่เพื่อ agglutinate กับ HCG coated red cells ซึ่งปฏิกิริยาดังกล่าวจะเกิดขึ้นประมาณ ๒ ชั่วโมงหลังการทดสอบ

\* แผนกสูติ-นรีเวชวิทยา คณะแพทยศาสตร์จุฬาลงกรณ์มหาวิทยาลัย